DISTANCES FORBIDDEN BY SOME TWO-COLORING OF \mathbb{Q}^2

MICHAEL REID, DOUGLAS S. JUNGREIS AND DAVE WITTE

Harvard University, Harvard University, and Massachusetts Institute of Technology

Received 6 October 1986, Revised 15 March 1988

0. INTRODUCTION

In 1973, D. R. Woodall [3] showed that there is a 2-coloring of \mathbb{Q}^2 (the rational points in the plane) that forbids the distance 1 (i.e. no two points of the same color are at distance 1 apart). Recently, this result was strengthened by P. D. Johnson, Jr. [2], who showed there is a 2-coloring of \mathbb{Q}^2 that simultaneously forbids all Euclidean distances of the form $\sqrt{p/q}$ with p and q odd positive integers. Johnson's 2-coloring obviously forbids certain other distances (e.g. $\sqrt{6}$) because no two rational points in the plane are precisely that far apart. We show in Section 1 that Johnson's coloring is optimal: for any distance d not covered by either of these two cases, no 2-coloring of \mathbb{Q}^2 forbids both of the distances 1 and d (see Theorem 1.5). This settles Problem 4 of [2]. In Section 2, we give a similar solution of Problems 2 and 3 of [2].

1. Two-colorings of \mathbb{Q}^2

Definition. Let $D \subseteq \mathbb{R}^+$. A 2-coloring of \mathbb{Q}^2 forbids the distances D if for all $x, y \in \mathbb{Q}^2$, with $d(x, y) \in D$, the two points x and y have different colors.

Definition. For any $D \subseteq \mathbb{R}^+$, we say there is an *odd D*-cycle (in \mathbb{Q}^2) if there exist $s_1, s_2, \ldots, s_n \in \mathbb{Q}^2$ such that $s_1 + s_2 + \cdots + s_n = 0$ (under componentwise addition), $||s_i|| \in D$ for each *i*, and *n* is odd. (Not having an odd *D*-cycle means a certain subset of \mathbb{Q}^2 is "weakly two-free" in the language of [1, 2].)

The following lemma follows from the fact that a graph is bipartite iff it contains no odd cycle.

¹⁹⁹¹ Mathematics Subject Classification. 05C15 (51K05).

Lemma 1.1. (cf. [1]) Let $D \subseteq \mathbb{R}^+$. There is no 2-coloring of \mathbb{Q}^2 that forbids D iff there is an odd D-cycle.

Lemma 1.2. (cf. [2, Lemma 1]) Let $d \in \mathbb{R}^+$ and set d' = kd, where k is any odd positive integer. Then there is an odd $\{1, d\}$ -cycle iff there is an odd $\{1, d'\}$ -cycle.

Proof. (\Rightarrow) Suppose $u_1 + u_2 + \cdots + u_l + s_1 + s_2 + \cdots + s_m = 0$, with l + m odd, and where $||u_i|| = 1$ for each *i*, and $||s_j|| = d$ for each *j*. Setting $s'_j = k \cdot s_j$, we have

$$k \cdot u_1 + k \cdot u_2 + \dots + k \cdot u_l + s'_1 + s'_2 + \dots + s'_m = 0,$$

and kl + m is odd.

(\Leftarrow) Suppose $u_1 + u_2 + \dots + u_l + s'_1 + s'_2 + \dots + s'_m = 0$. Setting $s_j = \frac{1}{k} \cdot s'_j$, we have $u_1 + u_2 + \dots + u_l + k \cdot s_1 + k \cdot s_2 + \dots + k \cdot s_m = 0$,

and l + km is odd.

Proposition 1.3. Let $p, q \in \mathbb{Z}^+$, with p even and q odd, and assume $d = \sqrt{p/q}$ occurs as a distance between rational points in the plane. Then there is an odd $\{1, d\}$ -cycle.

Proof. By Lemma 1.2 it suffices to show there is an odd $\{1, qd\}$ -cycle. Since d occurs as a distance, there are rational numbers α and β with $\alpha^2 + \beta^2 = pq$. Any integer that is a sum of two rational squares is a sum of two integer squares, so we may assume that α and β are non-negative integers. Now $\alpha \cdot (1, 0) + \beta \cdot (0, 1) + (-\alpha, -\beta) = 0$, and $||(-\alpha, -\beta)|| = \sqrt{pq} = qd$. Since $\alpha^2 + \beta^2 = pq$ is even, α and β have the same parity, so $\alpha + \beta + 1$ is odd.

Lemma 1.4. Let $D \subseteq \mathbb{R}^+$ and $d_0 \in \mathbb{R}^+$. Assume d_0 occurs as a distance between two rational points in the plane. Then there is an odd D-cycle iff there is an odd $(d_0 \cdot D)$ -cycle (where $d_0 \cdot D = \{d_0 d \mid d \in D\}$).

Proof. By assumption there are $\alpha, \beta \in \mathbb{Q}$ with $\sqrt{\alpha^2 + \beta^2} = d_0$. Then $\begin{pmatrix} \alpha & \beta \\ -\beta & \alpha \end{pmatrix} \in \mathrm{GL}_2(\mathbb{Q})$ is a linear transformation of the plane that preserves the set of rational points and expands all distances precisely by a factor of d_0 . The lemma follows.

Theorem 1.5. Let $d \in \mathbb{R}^+$. Assume there are $x, y \in \mathbb{Q}^2$ with d(x, y) = d, and that d can be written in the form $d = \sqrt{p/q}$ with $p, q \in \mathbb{Z}^+$ and p + q odd. Then no 2-coloring of \mathbb{Q}^2 forbids both of the distances 1 and d.

Proof. By Lemma 1.1, we need only show there is an odd $\{1, d\}$ -cycle in \mathbb{Q}^2 . By Lemma 1.4, there is an odd $\{1, d\}$ -cycle iff there is an odd $\{1/d, 1\}$ -cycle. So, replacing d by 1/d if necessary, we may assume $d = \sqrt{p/q}$ where p is even and q is odd. Then Proposition 1.3 applies.

Corollary 1.6. Let $D \subseteq \mathbb{R}^+$. No 2-coloring of \mathbb{Q}^2 forbids the distances D iff there are $d_1, d_2 \in D$ such that:

- (1) Each of d_1 and d_2 occurs as a distance between rational points in the plane; and
- (2) There exist $p, q \in \mathbb{Z}^+$ such that $d_1/d_2 = \sqrt{p/q}$ and p + q is odd.

2. Two-colorings of A_n and \mathbb{Q}^3

Let $A_n = \{(a_1/b_1, a_2/b_2, \dots, a_n/b_n) | a_i, b_i \in \mathbb{Z} \text{ and each } b_i \text{ is odd } \}$. We now prove the natural analogue of Theorem 1.5 for A_n and for \mathbb{Q}^3 . This settles problems 2 and 3 of [2].

Proposition 2.1. Let $p, q \in \mathbb{Z}^+$, with p even and q odd. Set $d = \sqrt{p/q}$ and assume there are $x, y \in A_n$ with d(x, y) = d. Then no 2-coloring of A_n forbids both of the distances 1 and d.

Proof. (cf. proof of Proposition 1.3) We need only show there is an odd $\{1, d\}$ -cycle in A_n (see Lemma 1.1). Write $d^2 = \alpha_1^2 + \alpha_2^2 + \cdots + \alpha_n^2$, where each α_i is rational with odd denominator. We may rescale in the manner of Lemma 1.2 to assume each α_i is an integer, so d^2 is an even integer. Then

$$\alpha_1 \cdot (1, 0, \dots, 0) + \alpha_2 \cdot (0, 1, \dots, 0) + \dots + \alpha_n \cdot (0, 0, \dots, 1) + (-\alpha_1, -\alpha_2, \dots, -\alpha_n) = 0,$$

and $\alpha_1 + \alpha_2 + \dots + \alpha_n + 1$ is odd.

Proposition 2.2. Let $d \in \mathbb{R}^+$. Assume there are $x, y \in \mathbb{Q}^3$ with d(x, y) = d, and that d can be written in the form $d = \sqrt{p/q}$ with $p, q \in \mathbb{Z}^+$ and p + q odd. Then no 2-coloring of \mathbb{Q}^3 forbids both of the distances 1 and d.

Proof. We need only show there is an odd $\{1, d\}$ -cycle in \mathbb{Q}^3 (see Lemma 1.1). Write $p/q = \alpha_1^2 + \alpha_2^2 + \alpha_3^2$ (with each $\alpha_i \in \mathbb{Q}$).

Case 1. p is even and q is odd. It follows that $\alpha_1, \alpha_2, \alpha_3$ have odd denominators (when written in lowest terms). Now apply the proof of Proposition 2.1.

Case 2. p is odd and q is even. At least one of $\alpha_1, \alpha_2, \alpha_3$ must have even denominator (when written in lowest terms). Assume it is $\alpha_1 = p_1/q_1$. Note

$$p_1 \cdot (1,0,0) + \frac{q_1}{2} \cdot (-\alpha_1, \alpha_2, \alpha_3) + \frac{q_1}{2} \cdot (-\alpha_1, -\alpha_2, -\alpha_3) = 0,$$

and $p_1 + q_1$ is odd.

Question. Is the analogue of Lemma 1.4 (and, hence, also the analogue of Corollary 1.6) true for A_n and \mathbb{Q}^3 ?

Acknowledgments

This work was done under the direction of Professor Joseph A. Gallian at the University of Minnesota, Duluth. Funding was provided by the Research at Undergraduate Institutions program of the National Science Foundation under Grant Number DMS8407498, with additional support from the University of Minnesota.

References

- 1. P. D. Johnson, Coloring abelian groups, Discrete Mathematics 40 (1982), 219-223, MR83k:20059.
- Peter D. Johnson, Jr., Two-colorings of a dense subgroup of Qⁿ that forbid many distances, Discrete Mathematics 79 (1990), no. 2, 191–195, MR91d:05046.
- D. R. Woodall, Distances realized by sets covering the plane, Journal of Combinatorial Theory, Series A 14 (1973), 187–200, MR46 #9868.